

| Serial # | Type | Grade | Size | Details |
|----------|----------|-------|------|---|
| 1 | DCIS | 2 | 0.20 | 0.2 DCIS largest focus |
| 2 | IDC/DCIS | 2 | n/a | invasive ductal carcinoma: well differentiated |
| 3 | IDC | 2 | 2.00 | nuclear grade 2 surrounding DCIS: cribriform type and intermediate grade 2 cm grade 2 |
| 4 | DCIS | 1 | 1.50 | 1.5 DCIS solid cribriform no necrosis identified grade 1 (low) |
| 5 | IDC/DCIS | 2 | 2.6 | RIGHT: 2.6 cm IDC; DCIS present comedo cribriform micropapillary solid grade 3; overall grade 2 |
| 6 | IDC/DCIS | 2 | 3 | LEFT: 3cm IDC; DCIS present cribriform solid grade 2; overall grade 2 |
| 7 | IDC | 1 | 1.00 | 1 cm invasive ductal grade 1 |
| 8 | DCIS | 3 | 1.4 | 1.4 cm DCIS solid comedo extensive comedonecrosis present grade 3 |
| 9 | DCIS | 3 | 0.00 | solid and cribriform types with punctate necrosis and nuclear grade 3 |
| 10 | DCIS | 1 | 0.6 | 0.6 cm Inv papillary cancer; cribriform DCIS grade 1 present; overall grade 1 |
| 11 | IDC/DCIS | 1 | 1 | 1 cm IDC; DCIS cribriform grade 1 present; overall grade 1 |
| 12 | IDC/DCIS | 3 | 1.1 | 1.1 cm IDC; DCIS comedo grade 3 present; overall grade 3 |
| 13 | IDC | 1 | 1.7 | 1.7 cm IDC; grade 1 |
| 14 | IDC/DCIS | 2 | 0.8 | 0.8 IDC; DCIS cribriform grade 2; overall grade 2 |
| 15 | DCIS | 3 | n/a | solid cribriform comedo DCIS grade 2/3 |
| 16 | DCIS | 2 | 0.6 | 0.6 cm solid cribriform grade 2 (intermediate) |
| 17 | PHY | n/a | n/a | excluded b/c phyllodes |
| 18 | IDC/DCIS | 1 | 1.5 | 1.5 cm IDC and DCIS solid cribriform with focal necrosis present; overall grade 1 |
| 19 | IDC | 1 | 1.8 | 1.8 cm; grade 1 |
| 20 | DCIS | 3 | n/a | grade 2 - 3 and cribriform |
| 21 | IDC/DCIS | 3 | 0.6 | 0.6 IDC; DCIS present comedo grade 3; overall grade 3 |
| 22 | DCIS | 3 | n/a | DCIS intermediate/high grade comedonecrosis |
| 23 | DCIS | 3 | n/a | high grade DCIS comedo type |
| 24 | IDC/DCIS | 2 | 0.4 | 0.4 cm IDC with DCIS grade 3 with expansive comedonecrosis present; overall grade 2 |
| 25 | DCIS | 3 | 4 | 4 cm DCIS solid grade 3 |
| 26 | DCIS | 2 | 0.5 | 0.5 cm DCIS cribriform grade 2 |
| 27 | DCIS | 2 | 0.1 | 0.1 cm DCIS micropapillary grade 2 |
| 28 | IDC | 3 | n/a | poorly differentiated |
| 29 | DCIS | n/a | 0.2 | 0.2 DCIS |
| 30 | IDC | 3 | n/a | INVASIVE, POORLY DIFFERENTIATED DUCTAL CARCINOMA. LYMPHATIC INVASION NOTED |
| 31 | IDC/DCIS | 1 | 1.7 | 1.7 cm IDC; DCIS of solid cribriform type grade 2; overall grade 1 |
| 32 | IDC | 1 | 0.2 | 0.2 IDC; no DCIS present; overall grade 1 |
| 33 | IDC | 2 | 0.9 | 0.9 cm and an additional 0.7cm nodule in posterior add margin; overall grade 2 (intermed); no DCIS present in final; |
| 34 | IDC/DCIS | 1 | 0.4 | invasive ductal 0.4 cm grade 1(low) necrosis not identified and cribriform DCIS present |
| 35 | IDC | n/a | n/a | |
| 36 | DCIS | 2 | 0.3 | 0.3 DCIS solid cribriform comedo type intermediate (2) grade |
| 37 | MUC/I | 1 | 0.1 | well diff mucinous <0.1cm; DCIS also seen cribriform and solid type with intermediate grade (2); overall grade 1 |
| 38 | DCIS | 3 | 0.5 | solid comedo DCIS HIGH GRADE (3); 0.5 cm extensive necrosis present |
| 39 | DCIS | 2 | n/a | nuclear grade 2 solid and cribriform |
| 40 | IDC/DCIS | 3 | 3.00 | INVASIVE DUCTAL CARCINOMA, POORLY DIFFERENTIATED, 3 CM SIZE. |
| 41 | IDC/DCIS | 2 | 3.9 | LYMPHOVASCULAR INVASION PRESENT |
| 42 | IDC | 3 | 2.7 | 3.9 IDC and solid DCIS present grade 3 (high); overall grade 2 |
| 43 | IDC/DCIS | 2 | n/a | INVASIVE DUCTAL CARCINOMA, 2.7 CM, GRADE 3. lymphovascular invasive identified |
| 44 | DCIS | 3 | n/a | IDC moderately diff with high grade DCIS with focal comedonecrosis |
| 45 | IDC | 2 | 0.25 | DUCTAL CARCINOMA IN SITU, NUCLEAR GRADE 3, SOLID SUBTYPE COMPRISES |
| 46 | IDC | 3 | 1.5 | LEFT: 0.25 intermediate grade (2) invasive ductal necrosis not identified 1.5 invasive pleomorphic carcinoma overall grade 3 |